



Listeriosis Phytotherapy: A Review Study on the Effectiveness of Iranian Medicinal Plants in Treatment of Listeriosis

Mahmoud Rafieian-Kopaei, PhD¹, Kourosh Saki, MD², Mahmoud Bahmani, PhD³, Sobhan Ghafourian, PhD⁴, Nourkhoda Sadeghifard, PhD⁴, and Morovat Taherikalani, PhD⁵

Abstract

Listeria monocytogenes can be found in many processed foods, raw milk, dairy products, meat and meat products such as sausages, beef and fish products, seafoods, eggs, fruits, and vegetables such as radish and cabbage. This article is a review study on the Iranian medicinal plants applied for treatment of listeriosis. Information of this review article was obtained by searching various key words such as *Listeria monocytogenes*, medicinal plants, plant extracts and essential oils among scientific articles published in databases of Google scholar, ISI Web of Knowledge, PubMed, Scopus, SID and Magiran. Thyme, German chamomile, great chamomile, yarrow, onion, oregano, nutmeg, sage, sagebrush, hyssop, rosemary, St John's wort, safflower, ajowan, cumin, peppermint, shallot, anise, and parsnip are known antilisteriosis medicinal plants. Bioactive phytochemicals, antioxidants and monoterpenes, sesquiterpene, coumarin, flavonoids, tannins, saponins, alkaloids, and terpenoids are the main ingredients of antilisteriosis medicinal plants.

Keywords

Listeria monocytogenes, medicinal plants, Iran

Received September 5, 2015. Received revised October 14, 2015. Accepted for publication October 27, 2015.

Listeria monocytogenes is a rod-shaped, Gram-positive, catalase-positive, oxidase-negative, and facultative anaerobic bacterium. Infections caused by this bacterium occur after consumption of contaminated water and food, and its symptoms are similar to flu, septicemia, and meningitis, especially in infants, pregnant women, the elderly, and immunocompromised individuals.¹

Listeria monocytogenes can be found in many processed foods, raw milk, dairy products, meat, and meat products such as sausages, beef and fish products, seafood, eggs, and in fruits and vegetables, such as radish, cabbage, and so on.²

Listeria monocytogenes grows in fish oil, smoked salmon, crab meat, cooked shrimp kept at -4°C , and canned lobster meat kept at 10°C .³ Food transmission is usually the main cause of sporadic and epidemic listeriosis in human. Recently, high rate of mortality and deaths of about 400 people in the United States due to listeriosis has become a matter of concern for food industries and regulatory organizations.⁴

Antimicrobial agents are chemical or natural compounds that are able to destroy or inhibit the growth of microscopic organisms.⁵ Several agents with animal, plant, or microbial sources play important protective role against microorganisms. The most important antimicrobial activity of compounds can be attributed to the antimicrobial enzymes (lactoperoxidase, lactoferrin), antimicrobial peptides (natural phenols), catechin hydroquinone, esters of fatty acids, phenolic antioxidants, antibiotics, and metals

(Cu).^{6,7} Medicinal plants play important roles in the treatment and prevention of diseases and improvement of population health in the communities.⁸⁻¹² Public interest in the use of herbal-based medications and natural products have been increasing especially in the recent years.¹³⁻¹⁶ Main reasons for this general approach are more side effects of chemical drugs and the environmental pollutions that threaten the earth.

Medicinal herbs as the effective and inexpensive sources of different drugs are used for the treatment and prevention of a wide variety of diseases, such as diabetes,¹⁷⁻¹⁹ liver

¹ Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

² Shahid Beheshti University of Medical Sciences, Tehran, Iran

³ Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran

⁴ Clinical Microbiology Research Center, Ilam University of Medical Sciences, Ilam, Iran

⁵ Razi Herbal Medicines Research Center & Department of Microbiology, School of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

Corresponding Author:

Morovat Taherikalani, Razi Herbal Medicines Research Center & Department of Microbiology, School of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran.

Email: taherikalani@gmail.com

Table 1. List of Native Iranian Medicinal Plants Against *Listeria monocytogenes*.

No.	Scientific Name	Family Name	Common Name	Result
1	<i>Zataria multiflora</i> Boiss	Lamiaceae	Thyme	Thymus vulgaris has antimicrobial effects on <i>Listeria monocytogenes</i> alone with MIC and MBC, 9.5 and 19 µg/mL, respectively. Combination of essential oil and nisin reduced the amount of MIC and MBC to 1.2 and 2.4 µg/mL, respectively. ⁴⁷
2	<i>Zataria multiflora</i> Boiss	Lamiaceae	Thyme	Nisin at both concentration of 500 and 1000 IU/g could not reduce <i>Listeria monocytogenes</i> count (<i>Listeria monocytogenes</i> PTCC 1163) to below the permissible limit for healthy people. Inhibitory activity of nisin against the growth of <i>Listeria monocytogenes</i> was reduced with time. While a combination of nisin and thyme oil at a concentration of 0.8% and 1.2% decreased number of bacteria below the permissible limit during 12 days. ⁴⁸
3	<i>Matricaria chamomilla</i> L	Asteraceae	German chamomile	MIC of German chamomile and great chamomile oils obtained in the range of 0.22-4 and 0.09-1 mg/mL, respectively. Differential inhibitory index of these plants essential oils demonstrated synergistic activity against <i>Listeria monocytogenes</i> , <i>Bacillus subtilis</i> , and <i>Bacillus cereus</i> and increased activity against <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , and <i>Salmonella typhimurium</i> . ⁴⁹
	<i>Tanacetum parthenium</i> L	Asteraceae	Great chamomile	
4	<i>Achillea eriophora</i> DC	Compositae	Yarrow	Yarrow's essential oil MIC against Gram-positive bacteria was 0.15-0.75 mg/mL and 1.5-3 mg/mL against Gram-negative bacteria, respectively. MIC against <i>Listeria monocytogenes</i> was 0.75 mg/mL. ⁵⁰
5	<i>Allium cepa</i> L.	Alliaceae	Onion	MIC for onion essential oil against <i>Listeria monocytogenes</i> was 125 µg/mL. ⁵¹ Results showed that antibacterial activity of oregano (with maximum and minimum growth inhibitory concentration = 0.62 µL/mL) was more effective than nutmeg (with minimal growth inhibitory concentration = 1.25 µL/mL and maximum growth inhibitory concentration = 2.5 µL/mL). ⁵²
6	<i>Origanum vulgare</i>	Labiatae	Oregano	
	<i>Myristica fragrans</i>	Myristicaceae	Nutmeg	
7	<i>Artemisia aucheri</i>	Asteraceae	Sage	MIC of aqueous extract of sage and sagebrush against <i>Listeria monocytogenes</i> was 80 mg/mL. ⁵³
	<i>Artemisia sieberi</i>	Asteraceae	Sagebrush	
	<i>Hyssopus officinalis</i> L	Lamiaceae	Hyssop	
8	<i>Rosmarinus officinalis</i>	Lamiaceae	Rosemary	The plants have been shown to be effective against bacteria in various doses and it had anti listeriosis in In-Vitro condition. ⁵⁴
	<i>Hypericum perforatum</i>	Hypericaceae	St John's wort	
	<i>Carthamus tinctorius</i>	Asteraceae	(Hardhay) Safflower	
9	<i>Carum copticum</i>	Umbelliferae	Ajowan	Ajowan essential oil has showed bacteriostatic effect at 0.15% and bactericidal effect at 0.3% concentration against <i>Listeria monocytogenes</i> in white fish extract. Adding 4% salt to white fish extract increased antibacterial activity of ajowan essential oil significantly. Growth of <i>Listeria monocytogenes</i> in the white fish extract reduced significantly in these concentrations (0.15% and 0.3%) compared with control. ⁵⁵
10	<i>Cuminum cyminum</i>	Apiaceae	Cumin	Count of <i>Listeria monocytogenes</i> in cheese samples containing 0.02% cumin essential oil showed 1 log reduction after 30 days and goes to zero after that. Presence of 0.04% essential oils caused bacterial reduction up to 1 log after 15 days, that goes to zero and was not isolated after that, while in control samples (cheese without essential oil) bacteria were isolated during all periods. ⁵⁶
11	<i>Mentha spicata</i>	Lamiaceae	Spearmint	MICs of peppermint essential oil and nisin were determined to be 160 and 320 µL/mL, respectively. ⁵⁷
12	<i>Ziziphora clinopodioides</i>	Lamiaceae	<i>Thymus vulgaris</i>	MIC of <i>Thymus vulgaris</i> for growth of <i>Listeria monocytogenes</i> was 1 µL/mL. ⁵⁸
13	<i>Allium ascalonicum</i>	Alliaceae	Shallot	
	<i>Pimpinella anisum</i>	Apiaceae	Anise	Results showed that shallot essential oil has highest antibacterial power against <i>Listeria monocytogenes</i> . Essential oil of anise showed good antibacterial effects on <i>Listeria monocytogenes</i> . ⁵⁹
14	<i>Parstinaca sativa</i>	Apiaceae	Parsnip	<i>Berberis vulgaris</i> has been shown to be effective against bacteria in various doses, however, <i>Daucus carota</i> on some doses. The activity of <i>Parstinaca sativa</i> , but not <i>Daucus carota</i> , was changed after 1 hour of keeping in refrigerator. Both types of carrots had significant inhibitory effects on <i>Listeria monocytogenes</i> but effect of parsnip was far more than carrots. ⁶⁰
	<i>Daucus carota</i>	Apiaceae	Carrot	

Abbreviations: MBC, minimum bactericidal concentration; MIC, minimum inhibitory concentration; PTCC, Persian type culture collection.

disorders,^{20,21} hyperlipidemia and related diseases,²²⁻²⁶ skin disorders,^{27,28} pain,²⁹⁻³² parasitic and bacterial diseases,³³⁻⁴⁰ respiratory diseases,⁴¹ anxiety, stress, and neurological diseases,⁴²⁻⁴⁴ and gastrointestinal diseases.^{45,46}

Methods and Materials

Information of this review article was obtained by searching various keywords such as *Listeria monocytogenes*, medicinal plant, plant extracts, and essential oils among scientific articles published in

databases such as Google Scholar, ISI Web of Knowledge, PubMed, Scopus, SID, and Magiran of which only publications in English and Persian languages published between 1976 and March 2015 were searched. Overall, 45 articles were retrieved, out of which 14 articles were excluded for being irrelevant to research purposes.

Results

Effective medicinal plants against *Listeria monocytogenes* are listed in Table 1.

Discussion

Recently antimicrobial effects of medicinal plants and their secondary metabolites have been studied.^{54,61} Chemical compositions of native medicinal plants in Iran that have positive antibacterial effects on *Listeria monocytogenes* have been reviewed in this study.

Carvacrol and thymol are the most prominent phenolic compounds with different levels in various parts of the thyme plant, including leaves, flowers, and roots. Thyme also contains tannins, flavonoids, saponins, and bitter substances.⁵⁴

German chamomile flowers extract contain flavonoids and coumarin. Kamazolen presence in chamomile gives blue color to its essential oil. Because of the existence of flavonoids, flowers of chamomile have a moisturizing and fragrance effect and for this reason are widely used in cosmetic industry.^{55,56} Camphor is the main chemical components of great chamomile essential oil.⁵⁷

Onion, because it contains flavonoids and alkyl-cysteine sulfoxide, has an important role in human health and nutrition. Alkyl-cysteine sulfoxide is the main precursor of onion flavoring materials.⁵⁸

Nutmeg essential oil contains borneol, geraniol, linalool, terpineol, eugenol, myristicin, and safrol.⁵⁹ Different parts of *Artemisia aucheri* plant comprise flavonoids, santonian, lipids, and bitter compounds.⁶⁰ Combinations of saffron Croatia, Christian, and crocin are picrocrocin.⁶² Rosemary is a rich source of phenolic compounds with antimicrobial effects against positive and negative bacteria.⁶³

Thymol, carvacrol, α - and β -pinene, parasimon compounds have been reported in ajowan essential oil.⁶⁴ Sabinene, flavonoids, polysaccharides, coumarin, cuminaldehyde, pinene, and terpinene are noted as the main components of the ajowan plant.⁶⁵

Chemical composition of the peppermint essential oil contains carvone, limonene, 1,8-cineole, linalool, menthol, menthone, and isomenthone.⁶⁶⁻⁶⁸

Allium genus plants such as shallot, garlic, and onions are rich in flavonols and organosulfur components.^{69,70}

Thymus vulgaris (*Ziziphora tenuior*) essential oil is composed of oxygen-containing monoterpene, pulegone, isomenthone, 1,8-cineole, and isopiperitone.⁷¹

Review of the obtained results of this study showed that bioactive phytochemicals, antioxidants and monoterpenes, sesquiterpene, coumarin, flavonoids, tannins, saponins,

alkaloids, and terpenoids are the main ingredients of antilisteriosis medicinal plants. Derivation of these compounds and complementary pharmacological studies can demonstrate their use as impressive drugs against *Listeria monocytogenes* and other pathogen bacteria. The mechanism by which these plants act against *Listeria* is still not clear; however, phenolic components of plants have been attributed to their antimicrobial activities. If this is true, other plants having these compounds⁷²⁻⁸² should also possess antibacterial activity and beneficial effect against this pathogen, which will be worth examining. Due to the to be increased of prevalence of antibiotic resistance in the bacterial diseases in Iran,⁸³⁻⁸⁶ identification of effective medicinal plants is necessary for finding natural active pharmaceutical ingredients for production of herbal antibiotics against the bacterial diseases. Should also possess antibacterial activity and beneficial effect against this pathogen, which will be worth examining.

Acknowledgments

The authors thank all people who helped in this study.

Author Contributions

All the authors wrote the first draft of the manuscript equally. MRK revised and edited the last version.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

Ethical Approval

Ethical issues (including plagiarism, misconduct, data fabrication, falsification, double publication or submission, redundancy) have been completely observed by the authors.

References

1. Murray PR, Baron EJ, Jorgensen JH, Pfaller MA, Tenover FC, Tenover FC. *Manual of Clinical Microbiology*. 7th ed. Washington, DC: American Society for Microbiology; 2003.
2. Gandhi M, Chikindas ML. *Listeria*: a foodborne pathogen that knows how to survive. *Int J Food Microbiol*. 2007;113:1-15.
3. Ben Embarek PK. Presence, detection, and growth of *Listeria monocytogenes* in seafoods. *Int J Food Microbiol*. 1994;23:17-34.
4. Hitchins AD. Detection and enumeration of *Listeria monocytogenes* in foods. In: Jackson GJ, Merker RI, Bandler R, eds. *Bacteriological Analytical Manual*. <http://www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm071400.htm>. Accessed November 27, 2015.
5. Anonymous. Antimicrobial agent. <http://www.answers.com/topic/antimicrobialagent>. 2009.
6. Appendini P, Hotchkiss JH. Review of antimicrobial food packaging. *Innov Food Sci Emerging Technol*. 2002;3:113-126.

7. Embuscado ME, Huber KC. *Edible Films and Coatings for Food Applications*. Springer, New York, NY: Springer; 2009:295-314.
8. Bahmani M, Mirhoseini M, Shirzad H, Sedighi M, Shahinfard N, Rafieian-Kopaei M. A Review on promising natural agents effective on hyperlipidemia. *J Evid Based Complementary Altern Med*. 2015;20:228-238. doi:10.1177/2156587214568457.
9. Bahmani M, Forouzan SH, Fazeli-Moghadam E, Rafieian-Kopaei M, Adineh A, Saberianpour SH. Oak (*Quercus branti*): an overview. *J Chem Pharm Res*. 2015;7:634-639.
10. Parsaei P, Karimi M, Asadi SY, Rafieian-Kopaei M. Bioactive components and preventive effect of green tea (*Camellia sinensis*) extract on post-laparotomy intra-abdominal adhesion in rats. *Int J Surg*. 2013;11:811-815. doi:10.1016/j.ijsu.2013.08.014.
11. Rabiei Z, Rafieian-Kopaei M, Heidarian E, Saghaei E, Mokhtari S. Effects of *Zizyphus jujube* extract on memory and learning impairment induced by bilateral electric lesions of the nucleus basalis of Meynert in rat. *Neurochem Res*. 2014;39:353-360. doi:10.1007/s11064-013-1232-8.
12. Bahmani M, Saki K, Golshahi H, et al. Ethnobotanical and therapeutic uses of camomille. *J Chem Pharm Res*. 2015;7:640-645.
13. Bahmani M, Rafieian-Kopaei M, Saki K, et al. Identification of medical plants acting on reproductive system disorders: an ethnobotanical study in Urmia, northwest of Iran. *J Chem Pharm Res*. 2015;7:493-502.
14. Delfan B, Kazemeini HR, Bahmani M. Identifying effective medicinal plants for cold in Lorestan Province, west of Iran. *J Evid Based Complementary Altern Med*. 2015;20:173-179. doi:10.1177/2156587214568458
15. Delfan B, Bahmani M, Hassanzadazar H, et al. Ethnobotany study of effective medicinal plants on gastric problems in Lorestan province, west of Iran. *J Chem Pharm Res*. 2015;7:483-492.
16. Bahmani M, Eftekhari Z, Jelodari Z, et al. Effect of Iranian herbal medicines in dysmenorrhea phytotherapy. *J Chem Pharm Res*. 2015;7:519-526.
17. Kafash-Farkhad N, Asadi-Samani M, Rafieian-Kopaei M. A review on phytochemistry and pharmacological effects of *Pran-gos ferulacea* (L.) Lindl. *Life Sci J*. 2013;10:360-367.
18. Bahmani M, Zargarani A, Rafieian-Kopaei M, Saki K. Ethnobotanical study of medicinal plants used in the management of diabetes mellitus in the Urmia, northwest Iran. *Asian Pac J Trop Med*. 2014;7(suppl 1):S348-S354.
19. Mirhoseini M, Baradaran A, Rafieian-Kopaei M. Medicinal plants, diabetes mellitus and urgent needs. *J Herb Med Pharmacol*. 2013;2:53-54.
20. Taghikhani A, Afrough H, Ansari-Samani R, Shahinfard N, Rafieian-Kopaei M. Assessing the toxic effects of hydroalcoholic extract of *Stachys lavandulifolia* Vahl on rat's liver. *Bratisl Lek Listy*. 2014;115:121-124.
21. Heidarian E, Rafieian-Kopaei M. Protective effect of artichoke (*Cynara scolymus*) leaf extract against lead toxicity in rat. *Pharm Biol*. 2013;51:1104-1109. doi:10.3109/13880209.2013.777931.
22. Rafieian-Kopaei M, Shahinfard N, Rouhi-Boroujeni H, Gharipour M, Darvishzadeh-Boroujeni P. Effects of *Ferulago angulata* extract on serum lipids and lipid peroxidation. *Evid Based Complement Alternat Med*. 2014;2014:680856. doi:10.1155/2014/680856.
23. Asgary S, Sahebkar A, Afshani M, Keshvari M, Haghjooyjavanmard Sh, Rafieian-Kopaei M. Clinical evaluation of blood pressure lowering, endothelial function improving, hypolipidemic and anti-inflammatory effects of pomegranate juice in hypertensive subjects. *Phytother Res*. 2014;28:193-199. doi:10.1002/ptr.4977.
24. Mirhoseini M, Baradaran A, Rafieian-Kopaei M. *Anethum graveolens* and hyperlipidemia: a randomized clinical trial. *J Res Med Sci*. 2014;19:758-761.
25. Khosravi-Boroujeni H, Mohammadifard N, Sarrafzadegan N, et al. Potato consumption and cardiovascular disease risk factors among Iranian population. *Int J Food Sci Nutr*. 2012;63:913-920.
26. Asgary S, Kelishadi R, Rafieian-Kopaei M, Najafi S, Najafi M, Sahebkar A. Investigation of the lipid-modifying and anti-inflammatory effects of *Cornus mas* L. supplementation on dyslipidemic children and adolescents. *Pediatr Cardiol*. 2013;34:1729-1735. doi:10.1007/s00246-013-0693-5.
27. Delfan B, Bahmani M, Eftekhari Z, Jelodari M, Saki K, Mohammadi T. Effective herbs on the wound and skin disorders: a ethnobotanical study in Lorestan province, west of Iran. *Asian Pac J Trop Dis*. 2014;4(suppl 2):938-942.
28. Asadi SY, Parsaei P, Karimi M, et al. Effect of green tea (*Camellia sinensis*) extract on healing process of surgical wounds in rat. *Int J Surg*. 2013;11:332-337. doi:10.1016/j.ijsu.2013.02.014.
29. Delfan B, Bahmani M, Rafieian-Kopaei M, Delfan M, Saki K. A review study on ethnobotanical study of medicinal plants used in relief of toothache in Lorestan Province, Iran. *Asian Pac J Trop Dis*. 2014;4(suppl 2):879-884.
30. Bahmani M, Shirzad HA, Majlesi M, Shahinfard N, Rafieian-Kopaei M. A review study on analgesic applications of Iranian medicinal plants. *Asian Pac J Trop Med*. 2014;7(suppl 1):43-53.
31. Rabiei Z, Rafieian-Kopaei M, Mokhtari S, Alibabaei Z, Shahrani M. The effect of pretreatment with different doses of *Lavandula officinalis* ethanolic extract on memory, learning and nociception. *Biomed Aging Pathol*. 2014;4:71-76.
32. Delfan B, Bahmani M, Hassanzadazar H, Saki K, Rafieian-Kopaei M. Identification of medicinal plants affecting on headaches and migraines in Lorestan Province, West of Iran. *Asian Pac J Trop Med*. 2014;7(suppl 1):376-379.
33. Karamati SA, Hassanzadazar H, Bahmani M, Rafieian-Kopaei M. Herbal and chemical drugs effective on malaria. *Asian Pac J Trop Dis*. 2014;4(suppl 2):599-601.
34. Bahmani M, Rafieian-Kopaei M, Hassanzadazar H, Saki K, Karamati SA, Delfan B. A review on most important herbal and synthetic antihelmintic drugs. *Asian Pac J Trop Med*. 2014;7(suppl 1):29-33.
35. Bahmani M, Saki K, Rafieian-Kopaei M, Karamati SA, Eftekhari Z, Jelodari M. The most common herbal medicines affecting *Sarcomastigophora* branches: a review study. *Asian Pac J Trop Med*. 2014;7(suppl 1):14-21.
36. Bahmani M, Karamati SA, Hassanzadazar H, et al. Ethnobotanic study of medicinal plants in Urmia city: identification and traditional using of antiparasites plants. *Asian Pac J Trop Dis*. 2014;4(suppl 2):906-910.

37. Bahmani M, Banihabib E, Rafieian-Kopaei M, Gholami-Ahangaran M. Comparison of disinfection activities of nicotine with copper sulphate in water containing *Limnatis nilotica*. *Kafkas Univ Vet Fak Derg*. 2015;21: 9-11.
38. Bahmani M, Farkhondeh T, Sadighara P. The anti-parasitic effects of *Nicotina tabacum* on leeches. *Comp Clin Pathol*. 2012;21: 357-359.
39. Bahmani M, Karamati SA, Banihabib EK, Saki K. Comparison of effect of nicotine and levamisole and ivermectin on mortality of leech. *Asian Pac J Trop Dis*. 2014;4(suppl 1):477-480.
40. Bahmani M, Saki K, Gholami-Ahangaran M, Parsaei P, Mohsenzadegan A, Zia-Jahromi N. Evaluating the anti-leech activity of methanolic extract of *Matricaria chamomilla* L. Comparing with ivermectin, mebendasole, praziquantel, rafoxanide, febantel and albendasole. *Mid East J Sci Res*. 2012;12:260-263.
41. Asadbeigi M, Mohammadi T, Rafieian-Kopaei M, Saki K, Bahmani M, Delfan B. Traditional effects of medicinal plants in the treatment of respiratory diseases and disorders: an ethnobotanical study in the Urmia. *Asian Pac J Trop Med*. 2014;7(suppl 1): 364-368.
42. Delfan B, Bahmani M, Hassanzadazar H, Saki K, Rafieian-Kopaei M. Identification of medicinal plants affecting on headaches and migraines in Lorestan Province, West of Iran. *Asian Pac J Trop Med*. 2014;7(suppl 1):376-379.
43. Saki K, Bahmani M, Rafieian-Kopaei M. The effect of most important medicinal plants on two important psychiatric disorders (anxiety and depression)-a review. *Asian Pac J Trop Med*. 2014; 7(suppl 1):34-42.
44. Saki K, Bahmani M, Rafieian-Kopaei M, et al. The most common native medicinal plants used for psychiatric and neurological disorders in Urmia city, northwest of Iran. *Asian Pac J Trop Dis*. 2014;4(suppl 2):895-901.
45. Hosseini-Asl K., Rafieian-Kopaei M. Can patients with active duodenal ulcer fast Ramadan? *Am J Gastroenterol*. 2002;97: 2471-2472.
46. Bahmani M, Zargar A, Rafieian-Kopaei M. Identification of medicinal plants of Urmia for treatment of gastrointestinal disorders. *Rev Bras Farmacogn*. 2014;24:468-480.
47. Rabiei S, Hosseini H, Rezaei M, Mousavi T. Inhibitory effects of Ajowan essential oil on growth of *Listeria monocytogenes* in Rutilus frissi kutum broth medium and fillet. *Iran J Nutr Sci Food Technol*. 2013;8:72-80.
48. Fazlara A, Sadeghi E, Rostami Soleimani P. Study on the antibacterial effects of *Cuminum cyminum* essential oil on *Listeria monocytogenes* in Iranian white cheese. *Iran J Food Sci Technol*. 2012; 9(35):35-44.
49. Mousavai MH, Mahmoudi R, Davoudi S, Shaveisi N. Antimicrobial effect of mint and niacin as synthetic form on *Listeria monocytogenesis*. *Med Plant J*. 2013;12:104-116.
50. Soltaninejad Sh, Setaei Mokhtari T, Rahbarian P. Antibacterial activity of essential oil and methanolic extract of *Zizifora cliniodis* on some pathogenic bacteria. *Microb Biotechnol Islamic Azad Univ*. 2010;2(5):1-6.
51. Hsani A, Mahmoudi R, Zare P, Hasany A. Biochemical properties and antimicrobial effects of *Allium ascalonicum* and *Pimpinella anisum* essential oils against *Listeria monocytogenes* in white brined cheese. *J Food Technol Res*. 2011;21:318-328.
52. Haghi Sayedjavadi N, Volasi N, Gamshidian M, Akhtari L. Comparison of *Parcatia sativa* and *Daucus carota* on *Listeria monocytogenes*. *Shahid Beheshti Univ Med J*. 2009;14:9-14.
53. Tepe B, Donmez E, Unlu M, Candan F, Daferera D, Vardar-Unlu G. Antimicrobial and antioxidative activities of the essential oils and methanol extracts of *Salvia cryptantha* (Montbret et Aucher ex Benth.) and *Salvia multicaulis* (Vahl). *Food Chem*. 2004;84: 519-525.
54. Burt S. Essential oils: their antibacterial properties and potential application in foods—a review. *Int J Food Microbiol*. 2004;94: 223-253.
55. Salamon I. Effect of the internal and external factors on yield and qualitative-quantitative characteristics of chamomile essential oil. *J Sci Ind Res*. 2007;99:132-139.
56. Franze C, Hoelzel J, Voemel A. Preliminary morphological and chemical characterization of some population and varieties of *Matricaria chamomila* L. *Acta Hort*. 2011;2:379-385.
57. Tiuman TS, Ueda-Nakamura T, Garcia Cortez DA, et al. Antileishmanial activity of parthenolide, a sesquiterpene lactone isolated from *Tanacetum parthenium*. *J Antimicrob Agent Chemother*. 2005;49:176-182.
58. Griffiths G, Trueman L, Crowther T, Thomas B, Smith B. Onions—a global benefit to health. *Phytother Res*. 2002;16: 603-615.
59. Tainter DR, Grenis AT. *Spices and Seasoning*. 2nd edition. New York, NY. Wiley; 2001.
60. Dinani NJ, Asgary A, Madani H, Naderi G, Mahzoni P. Hypocholesterolemic and antiatherosclerotic effect of *Artemisia aucheri* in hypercholesterolemic rabbits. *Pak J Pharm Sci*. 2010;23:321-325.
61. Rafieian-Kopaei M, Nasri H, Alizadeh F, Ataebi B, Baradaran A. Immunoglobulin A nephropathy and *Malaria falciparum* infection: a rare association. *Iran J Public Health*. 2013;42:529-533.
62. Abolhasani A, Bathaie SZ, Yavari I, Moosavimovahedi AA, Ghaffari M. Separation and purification of some components of Iranian saffron. *Asian J Chem*. 2005;17:727-729.
63. Tavassoli S, Mousavi SM, Emam-Djomeh Z, Razavi SH. Comparative study of the antimicrobial activity of *Rosmarinus officinalis* L. essential oil and methanolic extract. *Mid East J Sci Res*. 2011;9:467-471.
64. Aktug SE, Karapikar M. Inhibition of food borne pathogens by thymol, eugenol, menthol and ethanol. *Int J Food Microbiol*. 1987;4:161-166.
65. Demirci F, Guven K, Demirci B, Dadandi MY, Baser KH. Antibacterial activity of two *Phlomis* essential oils against food pathogens. *Food Control*. 2008;19:1159-1164.
66. Younis YMH, Beshir SM. Carvone-rich essential oils from *Mentha longifolia* (L.) Huds. ssp. *schimperii* Briq. and *Mentha spicata* L. grown in Sudan. *J Essential Oil Res*. 2004;16:539-541.
67. Zeinali H, Arzani A, Razmjoo R, Rezaee MB. Evaluation of oil compositions of Iranian Mints (*Mentha* spp.). *J Essential Oil Res*. 2005;17:156-159.
68. Shir-Mohammad H. *Generic and Herbal Medicines of Iran*. 2nd ed. Tehran, Iran: Arjomand Press; 2001.

69. Fattorusso E, Iorizzi M, Lanzotti V, Taglialatela-Scafati O. Chemical composition of shallot (*Allium ascalonicum* Hort.). *J Agr Food Chem*. 2002; 50 (20): 5686 -5690.
70. Mubarak AM, Kulatilleke CP. Sulfur constituents of need seed volatiles: a revision. *J Phytochem*. 1990;29:3351-3352.
71. Salehi P, Mirjalili MH, Hadian J, Nejad Ebrahimi S, Yousef Zadi M. Antibacterial activity and composition of the essential oil of *Ziziphora clinopodioides* subsp *bungeana* (JUZ.). Rech. F, from Iran Z. *Naturforsch*. 2006;61:677-680.
72. Bahmani M, Shirzad H, Rafieian S, Rafieian-Kopaei M. *Silybum marianum*: beyond hepatoprotection. *J Evid Based Complementary Altern Med*. 2015;20:292-301. doi:10.1177/2156587215571116.
73. Sharafati R, Sharafati F, Rafieian-Kopaei M. Biological characterization of Iranian walnut (*Juglans regia*) leaves. *Turk J Biol*. 2011;35:635-639.
74. Nasri H, Baradaran A, Ardalan MR, Mardani S, Momeni A, Rafieian-Kopaei M. Bright renoprotective properties of metformin: beyond blood glucose regulatory effects. *Iran J Kidney Dis*. 2013; 7:423-428.
75. Bahmani M, Saki K, Asadbeygi M, et al. The effects of nutritional and medicinal mastic herb (*Pistacia atlantica*). *J Chem Pharm Res*. 2015;7:646-653.
76. Nasri H, Nematbakhsh M, Rafieian-Kopaei M. Ethanol extract of garlic for attenuation of gentamicin-induced nephrotoxicity in Wistar rats. *Iran J Kidney Dis*. 2013;7:376-382.
77. Nasri H, Rafieian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci*. 2014;19:82-83.
78. Ghaed F, Rafieian-Kopaei M, Baradaran A, Nasri H. Ameliorative effects of metformin on renal histologic and biochemical alterations of gentamicin-induced renal toxicity in Wistar rats. *J Res Med Sci*. 2012;17:621-625.
79. Rafieian-Kopaei M, Nasri H. The ameliorative effect of *Zingiber officinale* in diabetic nephropathy. *Iran Red Crescent Med J*. 2014;16(5):e11324.
80. Nasri H, Rafieian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci*. 2014;19:82-83.
81. Baradaran A, Nasri H, Nematbakhsh M, Rafieian-Kopaei M. Antioxidant activity and preventive effect of aqueous leaf extract of Aloe vera on gentamicin-induced nephrotoxicity in male Wistar rats. *Clin Ter*. 2014;165:7-11. doi:10.7471/CT.2014.1653.
82. Nasri H, Rafieian-Kopaei M. Medicinal plants and antioxidants: why they are not always beneficial? *Iran J Public Health*. 2014; 43:255-257.
83. Taherikalani M, Etemadi G, Geliani KN, Fatollahzadeh B, Soroush S, Feizabadi MM. Emergence of multi and pan-drug resistance *Acinetobacter baumannii* carrying blaOXA-type -carbapenemase genes among burn patients in Tehran, Iran. *Saudi Med J*. 2008;29(4):623-624.
84. Pakzad I, Ghafourian S, Taherikalani M, Sadeghifard N, Abtahi H, Rahbar M, Mansory Jamshidi N. qnr Prevalence in Extended Spectrum Beta-lactamases (ESBLs) and None-ESBLs Producing *Escherichia coli* Isolated from Urinary Tract Infections in Central of Iran. *Iran J Basic Med Sci*. 2011;14(5):458-464.
85. Shamsavan S, Emameini M, Noorazar Khoshnab B, Khoramian B, Asadollahi P, Aligholi M, Jabalameli F, Eslampour MA, Taherikalani M. A high prevalence of mupirocin and macrolide resistance determinant among *Staphylococcus aureus* strains isolated from burnt patients. *Burns*. 2012;38(3):378-382.
86. Haghi-Ashteiiani M, Sadeghifard N, Abedini M, Soroush S, Taheri-Kalani M. Etiology and antibacterial resistance of bacterial urinary tract infections in Children's Medical Center, Tehran, Iran. *Acta Med Iran*. 2007;45(2):153-157.